

Sources of contamination in the emission from the Local Bubble



Anjali Gupta

Massimiliano Galeazzi

Eugenio Ursino



Philadelphia, 22 April 2008

Outline

- **The Diffuse X-ray background**
- **Unidentified point sources**
 - **Spectral Properties**
- **The Warm Hot Intergalactic Medium (WHIM)**
 - **Spectral properties**
 - **Angular properties**


Philadelphia, 22 April 2008

The Diffuse X-ray Background







Philadelphia, 22 April 2008


The Diffuse X-ray Background




Identification of the point sources



Mosaic images of the targets used in this investigation. The white circles represent the identified point sources.




The logN-logS



The averaged logN-logS for all the targets used in our investigation

Philadelphia, 22 April 2008


Spectral Properties of the point sources



*Average spectrum of all the sources detected in the **0.5-2.0 keV** (black circles) and **0.4-0.6 keV** (pink circles) bands.*


*The black and red curve represent the powerlaw fit with the photon index of **1.77 ± 0.01** and **1.93 ± 0.01** , for the two bands respectively.*

Spectral Properties of the bright and the faint sources



*Average spectrum of the bright (black) and faint (pink) sources detected in the **0.5-2.0 KeV**.*


*The black and red curve represent the powerlaw fit with the photon index of **1.87±0.01** and **1.40±0.05**, for the bright and faint sources respectively.*




*Average spectrum of the bright (black) and faint (pink) sources detected in the **0.4-0.6 KeV**.*

*The black and red curve represent the powerlaw fit with the photon index of **2.05±0.01** and **1.69±0.04**, for the bright and faint sources respectively.*

The Warm Hot Intergalactic Medium



Cen & Ostriker, 1999




Borgani et al. 2004



Spatial distribution of the WHIM at $z \sim 0$

Philadelphia, 22 April 2008

Flux vs. Redshift






*Expected average flux in the energy range **0.370-0.925 keV** due to the WHIM as a function of red-shift.*

*The total average flux due to the WHIM is about **5.7 photons/cm²/s/sr***





*The total DXB flux is about **29.3 photons/cm²/s/sr** (derived using data from McCammon et al., 2002 - ApJ 576, 188).*

Spectral Distribution of the WHIM




Philadelphia, 22 April 2008


WHIM signature in the Angular distribution of X-ray images



Mosaic images of the targets used in this investigation. The white circles represent the identified point sources.



Angular Autocorrelation




$$w(\theta) = \frac{\langle I(n)I(n') \rangle}{\langle I \rangle^2} - 1$$

The fraction of X-rays due to the WHIM in the energy band 0.4-0.6 keV is


$$18 \pm 5\%$$

of the total diffuse X-ray emission.



Calculated AcF for the two control targets, MBM20 (squares) and the Eridanus hole (full circles), compared with the average AcF (empty circles).

Conclusion




The End

Philadelphia, 22 April 2008

Angular Autocorrelation

- $$n^2 w(\Theta) = n_p^2 w_p(\Theta) + n_b^2 w_b(\Theta) + n_w^2 w_w(\Theta)$$



ACF of the diffuse X-ray background and its components.

Acf of Hubble Deep field North

